

New mineralisation windows in East Riverina: Results from the five-year mapping project

Contents



- Introduction
- Early Silurian epithermal–porphyry systems
- Early Devonian intrusion-related systems
 - Intrusion-related tin
 - Intrusion-related gold
 - Porphyry–epithermal gold
- Early to Middle Devonian structurally controlled low sulfidation gold mineralisation
- Conclusions

Introduction

East Riverina mapping project

5-year project (mapping 2014 to 2018)

- finalisation of line work, data and reports in 2019.

Update geological knowledge

- previous maps mostly from 1960–70s at 1:250 000
- increasing land use pressures.

Multi-disciplinary approach

- ‘Boots on the ground’ mapping with specialist input.

Applied research projects


- e.g. Uni of Newcastle, Lachlan Orogen ARC.

Local engagement



- professionals and community.



GOAL 1
**A growing and
diverse economy**
**Direction 12: Sustainably
manage mineral resources**



Surface geology from
NSW Seamless Geology
over 1vd TMI



Source: Glen 2013

New data

>7250 new observations & measurements (FieldObs)

- available in MinView
- includes photos, sample and analytical information.

~200 new age determinations


- >100 new isotopic:
 - SHRIMP (Geoscience Australia)
 - Ar–Ar (UoM, ANU), LA ICPMS (UoN, JCU, MU)
- >90 new palaeontological:
 - conodonts, graptolites, fish and invertebrates.



Australian Government
Geoscience Australia



FieldObs with surface geology
(NSW Seamless Geology over 1VD TMI).



New linework


Into Seamless Geology

Four provinces:


- Lachlan Orogen (LAO)
- Western Devonian Basins (WDB)
- Cenozoic Igneous Province (CIP)
- Cenozoic Sedimentary Province (CSP).



Surface Geology (all provinces)
(NSW Seamless Geology over 1VD TMI).



LAO and WDB
(NSW Seamless Geology over 1VD TMI).




New ideas

Available via East Riverina webpage:

- 18 x GS reports – field work
- 2 x GS reports – palaeontology
- 6 x GS reports – SHRIMP U–Pb (co-branded as Geoscience Aust. records)
- 2 x GS reports – LA ICPMS U–Pb
- 1 x GS report – mineral systems
- 1 x GS report – hydrogeochemistry (CSIRO)
- GS report, QN, paper – remote sensing (RMIT).



	Miners & explorers	Investors	Landholders & community	Resources Regulator			
IMER	Geoscience information	Applications & approvals	Safety and health	Compliance & reporting	Programs & initiatives	Rules & forms	Fossicking in NSW

[Home](#) / [Miners & explorers](#) / [Geoscience information](#) / [Projects](#) / [East Riverina Mapping Project](#)

Geological Survey of NSW

NSW State Emblems

NSW geology overview

Projects

Services

Products and data

Geotrails

Key contacts

MinEx CRC

East Riverina Mapping Project



The East Riverina Mapping Project provides up-to-date data to aid mineral exploration and to inform regional land-use decisions. Land-use pressure has increased in the east Riverina region, particularly in the new residential zones around Wagga Wagga and along the Hume Highway corridor. The latest Wagga Wagga 1:250 000 geological map was released in 1966 before the development of modern mapping and data collection techniques that allow interpretation of geology below deep alluvial cover.

The project area is in the central Lachlan Orogen and includes the Wagga tin or metamorphic belt, and the Gilmore and Kancoona fault zones. The area is prospective for a range of deposit-styles, including intrusive tin-tungsten deposits (e.g. Ardlethan), orogenic gold and copper-gold porphyry systems.



Systematic field mapping is integrated with specialist studies such as mineral systems, geochronology (with Geoscience Australia), palaeontology, spectral analysis, geophysical modelling, reinterpretation of seismic lines, 3D modelling of geology and fluid flow, and hydrogeochemistry (with CSIRO).

Products will include 1:100 000 (or better) geological mapping that will be incorporated into statewide digital geoscience datasets, annual technical and specialist reports, mineral prospectivity assessments and 3D models.

- 2 x Masters thesis, 2 x Honours thesis (UoN)
- Various conference posters, abstracts, talks
- Plus more reports coming soon
 - e.g. mineral systems and synthesis reports.

Today's focus

Mineral systems associated with Siluro-Devonian magmatism and deformation



Early Silurian epithermal–porphyry systems

Epithermal–porphyry systems


Magnetic linears along Gilmore Fault Zone
associated with intermediate to mafic rocks

Previously interpreted as:

- Continuation of the Junee–Narromine Volcanic Belt of the Macquarie Igneous Province (Benambran Cycle);
- Associated with opening of the Tumut Trough (Tabberabberan Cycle);
- Of uncertain or mixed origin.




LAO geology (ERIV)
over 1VD TMI.




Epithermal–porphyry systems

Gidginbung–Temora belt

- Part of Junee–Narromine Volcanic Belt (MIP)
- Late Ordovician – Early Silurian
- Epithermal mineralisation
 - e.g. **Gidginbung**
- Porphyry mineralisation
 - e.g. **Donnington**



Source: Sandfire Resources, Cronin et al. (2017)




LAO geology (ERIV)
over 1VD TMI.

Epithermal–porphyry systems

Junawarra Volcanics

- Associated with high sulfidation epithermal and carbonate base-metal mineralisation at Dobroyde (Diemar and Hughes 2018).
- First U–Pb geochronology
 - 431.1 ± 1.8 Ma for a porphyritic andesite at Dobroyde
 - 432 Ma (prelim.) for an autoclastic andesitic breccia to south.
- Tholeiitic affinity, with trace element characteristics indicative of a non-subduction setting (Eastlake 2018).
- New mapping of ‘jasper’ overlying mafic rocks
 - evidence of hydrothermal fluids associated with eruption.






LAO geology (ERIV)
over 1VD TMI.

Source: Waltenberg et al. 2019, Bodorkos et al. in prep.,
ERIVKSH0112, ERIVMAE1221, Eastlake (2018)
GS2018/0255.

Epithermal–porphyry systems

Cooba monzonite

- High-K, subduction-related monzonite identified
 - 439 Ma (prelim. age, Bodorkos et al. in prep.)
- Similar age and geochemical affinity to mineralised intrusions at Cadia and Goonumbla
- Associated with historical copper workings.




Epithermal–porphyry systems

Implications:

- Junawarra Volcanics younger (~432 Ma) with tholeiitic geochemical affinity
 - Represent opening of Tumut Trough
 - Contemporaneous with S-type granites.
 - Associated with Dobroyde and hydrothermal activity
- Monzonite at Cooba has similar age and geochemistry to Cadia and Goonumbla
 - Extends ~439 Ma Phase 4 Macquarie Igneous Province to south.



Source: Eastlake (2017) after Crawford et al. (2007)



Early Devonian intrusion-related systems

Intrusion-related tin

The southern part of the 'Wagga Tin Belt'.


Largest resource in ERIV at Ardlethan with hard-rock mining (1965–1986) until world tin price collapsed.

- Unmined total resource (tailings, waste, hard rock) of 66,500 t of contained tin.

Also tin–tungsten at Burrandana.





Surface geology (NSW Seamless Geology)
over 1VD TMI.



Intrusion-related tin

Ardlethan area


- Breccia pipes (e.g. Ardlethan mine)
- Tin-bearing greisen zones and veins (e.g. Bygoo).



Intrusion-related tin

What age is mineralisation?

- Previously assumed to relate to the S-type, ~430 Ma Koetong Supersuite
- Though Ardlethan thought to be younger (Blevin 2010)
- New mapping, petrography and geochronology have differentiated magmatic history.
- Tin mineralisation is younger
 - ~414 Ma.




Source: Bull & Blevin (2016)

Intrusion-related tin

Implications:

- Tin prospectivity study completed pre-East Riverina mapping project
 - ~430 Ma Koetong Supersuite was a key theme in weights of evidence approach
 - Now know that tin mineralisation is associated with Early Devonian magmatism (~414 Ma)
- Upcoming exploration model and mineral potential mapping for intrusion-related tin in zone 55 west
 - Will incorporate new data and ideas
- Mapping has constrained exploration search space.



Intrusion-related gold

Hobbs Pipe (near Mount Adrah) focus

- 20.1 Mt @ 1.12 g/t Au for 727,727 oz Au.

Several quartz monzonite to monzodiorite intrusions identified in Ordovician Abercrombie Formation country rock

- Hobbs Pipe dated at 414.7 ± 2.6 Ma, SHRIMP U–Pb zircon (Waltenberg et al. 2015).


Gold mineralisation:

- Disseminated within Hobbs Pipe intrusive body
- Quartz reefs in Abercrombie Formation.



Source: White (2013), 0.50 g/t cut off

Surface geology (NSW Seamless Geology)
over 1VD TMI.



Intrusion-related gold

New GS report by Wang et al. (in prep.)

- Integration of HyLogger™ data, petrography, sulfur isotopes
- Confirms intrusion-related gold genesis.



Sericite–calcite–leucoxene–pyrite–arsenopyrite alteration:

- Associated with gold mineralisation in the intrusion
- Within margins of high-grade gold-bearing reefs in host rocks
- Overprints regional foliation.

Sulfur isotopes indicated magmatic-dominated source.



Source: Waltenberg et al. 2015, Wang et al. (in prep.)



Intrusion-related gold

New GS report by Wang et al. (in prep.)

- Integration of HyLogger™ data, petrography, sulfur isotopes
- Confirms intrusion-related gold genesis.



Sericite–calcite–leucoxene–pyrite–arsenopyrite alteration:

- Associated with gold mineralisation in the intrusion
- Within margins of high-grade gold-bearing reefs in host rocks
- Overprints regional foliation.

Sulfur isotopes indicated magmatic-dominated source.



Source: Waltenberg et al. 2015, Wang et al. (in prep.)




Intrusion-related gold

Implications:

- Intrusion driver for mineralisation
 - Intrusion
 - Country rock.
- Hobbs Pipe is exposed mineralised ~415 Ma intrusion in hanging wall of the Gilmore Fault Zone.
- Other blind intrusions in same structural setting?
 - Leucoxene as vector?



Source: Geoscience Australia,
Spampinato (2018) – GS2018/0576,




Porphyry–epithermal gold

Focus on the Traist prospect in Howlong area



Surface geology
(NSW Seamless Geology) over 1VD TMI.




Porphyry–epithermal gold

Porphyritic intrusions along a splay of the Kancoona Fault.

- Gold mineralisation hosted by porphyritic intrusions and Ordovician Abercrombie Formation country rock.
- Dextral strike-slip movement interpreted from aeromagnetic imagery
- Consistent with kinematic history of Kancoona–Kiewa fault zone in Victoria.

New mapping, petrography and geochronology for Jindera Granite.

- Series of metaluminous, highly fractionated, I-type lobes younging to the south (Bull 2017)
- ~407 to ~403 Ma (Bodorkos et al. 2015, in prep.).




Porphyry–epithermal gold

Porphyritic intrusions along a splay of the Kancoona Fault.

- Gold mineralisation hosted by porphyritic intrusions and Ordovician Abercrombie Formation country rock.
- Dextral strike-slip movement interpreted from aeromagnetic imagery
- Consistent with kinematic history of Kancoona–Kiewa fault zone in Victoria.

New mapping, petrography and geochronology for Jindera Granite.


- Series of metaluminous, highly fractionated, I-type lobes younging to the south (Bull 2017)
- ~407 to ~403 Ma (Bodorkos et al. 2015, in prep.).



Porphyry–epithermal gold

New GS report by Forster et al. (in prep.):

- Integration of HyLogger™ data, petrography, geochronology, sulfur and lead isotopes
- New Re–Os age of molybdenite of 404.4 ± 1.5 Ma at Traist – within error of the Jindera Granite
- S isotopes – magmatic-dominated sulfur source
- Pb isotopes – typical Lachlan Orogen Early Devonian granite signature
- Concludes that mineralisation at Traist is epithermal carbonate base-metal–gold mineralisation.




Porphyry–epithermal gold

Implications:

- Newly recognised ~405 Ma magmatic event with gold mineralisation
- Intrusions along splays of major fault system
 - Kancoona (Boothagandra)–Kiewa FS
- Poor exposure
 - Traist porphyry is only intrusion exposed.



Surface geology
(NSW Seamless Geology) over 1VD TMI.



Early to Middle Devonian structurally controlled low sulfidation gold mineralisation

Structurally controlled gold

Gold mineralisation along Gilmore Fault Zone


- Plus parallel faults and related faults.

New GS report (Stuart et al. in prep.) reviewed individual goldfields in a regional context

- >500 occurrences with 32.2 t (1.14 Moz) Au
- Gold is hosted by range of Ord–Sil–Dev lithologies
- Timing Early (Bindian) to Middle Devonian (Tabberabberan) – though little direct dating.




Gold occurrences (MetIndEx) with surface geology
(NSW Seamless Geology) over 1VD TMI.



Structurally controlled gold

Three main settings ...

1. Narrow vein arrays within relatively competent rock types adjacent to second or third order splays of the GFZ and parallel structures.
 - e.g. Adelong, West Wyalong.




Source: Stuart and Ricketts (2019) from Stuart et al. (in prep.). After Woodcock & Fischer (1986).

Gold occurrences (MetIndEx) with LAO geology (NSW Seamless Geology) over 1VD TMI.

Structurally controlled gold

Three main settings ...

2. Along contacts between granites and metasedimentary country rock, or in pressure shadows near granites, where fluids have been focussed along contacts or around competent bodies.
 - e.g. Grong Grong.



Source: Stuart and Ricketts (2019) from Stuart et al. (in prep.). After Woodcock & Fischer (1986).



Gold occurrences (MetIndEx) with LAO geology (NSW Seamless Geology) over 1VD TMI.

Structurally controlled gold

Three main settings ...

3. Gold associated with folding adjacent to the GFZ.

- e.g. Barmedman.




Source: Stuart and Ricketts (2019) from Stuart et al. (in prep.) after Eastlake (2016), Lewis and Downes (2008)

Gold occurrences (MetIndEx) with LAO geology (NSW Seamless Geology) over 1VD TMI.

Structurally controlled gold

Implications:

- Structural setting key
 - Crustal scale and associated faults – geometry and kinematics
 - Lithology of host not as critical as inflections, contrasts, contacts etc.
- Multiple Devonian gold-forming events
- Importance of quality mapping to identify key criteria for mineral potential mapping.



Conclusions


Conclusions

New data, line work and ideas from mapping project

Increased understanding of:

- Siluro-Devonian magmatic history
- Structural framework, kinematics and history.

Implications for mineral systems and exploration



The work of many!

Geological Survey of NSW:

- Phil Blevin, Michael Bruce, Kate Bull, Lorraine Campbell, Astrid Carlton, Daniel Cronin, Liann Deyssing, Mark Eastlake, David Forster, John Greenfield, Kyle Hughes, Karen Montgomery, Bob Musgrave, Ian Percival, Glen Phillips, Mel Ricketts, Lawrie Sherwin, Cait Stuart, Steven Trigg, Yamei Wang, Brad Williams, Yong-Yi Zhen.

Geoscience Australia:

- Simon Bodorkos, Kathryn Waltenberg, Phil Main.

Uni of Newcastle:

- Michael Bell, Bill Collins, Alistair Hack, Deepika Venkataramani, David Boutelier.

Previous workers – exploration, government, academics





Phil Gilmore

phil.gilmore@planning.nsw.gov.au